BIRDING ALONG OUR CREEKS

Instructions: Use binoculars to identify birds. In a notebook, write down the species of birds you see–if known. If unknown, write down a detailed description with size, color, shape, beak, and behavior. **Please do not disturb nests or birds.**

Name:		Date:
Location	Mosthan	

Common Birds:


Female:♀ Male:♂



Canada Goose Branta canadensis

Very common and widespread. Found on or near any body of water. Often seen grazing on lawns, golf courses, and farmlands. Easily identified by long black neck.

Seen: 🗹



American Kestrel Falco sparverius

Common in non-breeding season (Aug-Mar). Often seen on tall wires and fence posts. The smallest falcon in North America. Identified by boldly patterned head.

Seen: 🗌



Great Blue Heron Ardea herodias

Uncommon. Forages in open habitats: grasslands and shallow water. Usually solitary and most active at night. Identified by large body and grayish-blue hue.


Seen: 🗌



Brewer's Blackbird

Euphagus cyanocephalus Very common and widespread. Often seen in parks, parking lots, and other open areas. Males are dark bluish-black with yellow eyes. Females: mostly brown.

Seen: 🗌



American Coot

Common. Often seen foraging in small ponds with plants near the surface or grazing on lawns. Forms large compact flocks. Duck-like with stubby-white bill.

Seen: 🗌



Black-billed Magpie Pica hudsonia

Very common and widespread. Often seen in urban areas, parks, and grasslands with scattered trees. Identified by black and white body and long black tail.

Seen: 🗌

BIRDING ALONG OUR CREEKS

Common Birds:



Killdeer Charadrius vociferus

Very common and widespread. Nests and forages in parks, parking lots, and other open areasoften far from water. Easily identified by two black stripes on throat.


Seen: 🗌



American Pelican

Pelecanus erythrorhynchos Common in breeding season (Apr-Jul). Often in flocks, soaring high overhead. Feeds by scooping small fish in bill pouch. Identified by large yellow bill.





Black-chinned Hummingbird Archilochus alexandri

Common in non-breeding season (Aug-Mar). Often seen in wooded areas. Males have black chin and purple throat.

Seen: 🗌



White-crowned Sparrow Zonotrichia leucophrys

Common in non-breeding season (Aug-Mar). Nests in brushy areas surrounded by open habitats, like parks.

Seen:

Notes:

Questions: What birds are here? How many are there? How are they using the space?

